

+

Physician's Orders GENERAL PRE-SCHEDULED SURGERY PRE-PROCEDURE

Surgeon/Physician	
Patient name	Date of birth
REQUIRED: Prepare consent (Consent to read)	
REQUIRED: Weightkilograms (only) REQUIRED: Alle	rgies
PENICILLIN ALLERGY? No Yes, reaction No anaphylaxis. May give Cephalosporin Anaphylaxis. No Cephalosporin	
REQUIRED (must choose one): A.M. Admit: Admit to Inpatient Outpatient Potential extended recovery (patient remains outpatient status, but may re	
CONSULTS: PHYSICIAN CONSULT: Reason: Medical clearance. Name	LABORATORY: All labs will be completed using Anesthesia Guidelines. Any orders checked below will be completed on Admission Day of Surgery, in
PT CARE/ACTIVITY: ☐ Hair clipping pre-procedure in Surgical Center as indicated. Site ☐ Chlorhexidine cloth (CHG) skin cleansing pre-procedure Site ☐ TEDs: ☐ Knee high ☐ Thigh high ☑ Pneumatic compression device: Comments: For surgical cases lasting greater than 60 minutes and/or high risk patients as defined by Anesthesia Guidelines. NOTE: If patient does not meet the Spectrum Health criteria above and you still want the device(s) placed, then SCD's must be ordered below. ☐ Knee high, bilaterally ☐ Knee high, right leg ☐ Knee high, left leg ☑ Protocol - pre-procedure anesthesia orders: Pre-procedure - May initiate pre-operative anesthesia orders (include laboratory and diagnostic tests)	addition to prior lab testing required by the Anesthesia Guidelines. Basic metabolic panel

Patient Name
DOB
MRN
Physician

CSN

 ${\bf Telephone\ order/Verbal\ order\ documented\ and\ read-back\ completed.\ \ Practitioner's\ initials\ _$

CONTINUED ON PAGE 2 →

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.

TRANSCRIBED:		VALIDATED:		ORDERED:		
TIME	DATE	TIME	DATE	TIME	DATE	Pager #
	Sign		R.N. Sign		Physician Print	Physician

CONFIDENTIAL NOTICE: The content of this fax is intended only for the named recipient(s) and may contain information that is protected under applicable law. If you are not the intended recipient(s) or if you receive this fax in error, notify the sender at the address or telephone number above. Also destroy any copies.

DO NOT MARK BELOW THIS LINE

BARCODE ZONE

DO NOT MARK BELOW THIS LINE





+

GENERAL PRE-SCHEDULED SURGERY - PRE-PROCEDURE (CONTINUED)

Page 2 of 2

ALL DESIRED ORDERS MUST BE CHECKED OR COMPLETED.

IV SOLUTIONS:
∠ Lidocaine 1% 0.25 to 2 mL ID for IV starts
☐ Lactated ringers solution 1000 mL IV, 100 mL/hour
\square 0.9% sodium chloride 1000 mL IV, 100 mL/hour
MEDICATIONS:
ANTIMICROBIALS (PROPHYLACTIC):
☐ Cefazolin 2 grams IV, if patient is less than 120 kg administered per anesthesia
☐ Cefazolin 3 grams IV, if patient is greater than or equal to 120 kg administered per anesthesia
PENICILLIN (PCN) ALLERGY:
FOR IMMEDIATE TYPE PENICILLIN ALLERGY:
☐ Clindamycin:
(when the lumen of the intestinal tract <u>IS NOT</u> entered)
• 900 mg IV administered per anesthesia
☐ Clindamycin PLUS Gentamicin:
(when the lumen of the intestinal tract <u>IS</u> entered)
 Clindamycin 900 mg IV administered per anesthesia Gentamicin 5 mg/kg IV administered per anesthesia
FOR CURRENT OR HISTORY OF MRSA COLONIZATION OR MRSA RISK FACTORS:
[residence in long-term healthcare facility (not assisted living), HD patient, tracheostomy or gastrostomy tube, chronic decubitus ulcer or wound, long term vascular access]:
Vancomycin (start administration within 120 minutes before
skin incision):
☐ 1 gram IV, if patient is less than 70 kg administered per anesthesia
☐ 1.5 grams IV, if patient is 70-100 kg administered per anesthesia
 2 grams IV, if patient is greater than 100 kg administered per anesthesia
PREPS:
☐ Enema: Type Time
LOCAL ANESTHESIA - HEMORRHOIDECTOMY PROCEDURES ONLY:
Use of Exparel is restricted to hemorrhoidectomy procedures only.
☐ Bupivacaine liposome (EXPAREL) 1.3% injection, 266 mg infiltration administered intra-operative

Patient Name			
DOB			
MRN			
Physician			
CSN			

MEDICATIONS: (CONTINUED)
VTE PROPHYLAXIS:
 Enoxaparin 40 mg subQ upon arrival NOTE: If spinal or epidural anesthesia, or peripheral nerve block
are planned, DO NOT use enoxaparin. Use subQ heparin instead.
☐ Heparin 5000 unit(s) subQ upon arrival
RESPIRATORY:
☐ Incentive spirometer
BETA BLOCKER:
Continue current therapy with sips of water in AM. Contact anesthesia for order if patient did not continue beta blocker
therapy
Medication
Dose Route Frequency
OTHER: NOTE: • For any additional orders: handwrite clearly or type below.
Must check the box for order to be processed.
<u> </u>
NOTE: • If there is a frequent order that needs to be added to your form,
contact Grand Rapids Spectrum Health Surgical Pre-procedure
Planning Manager or Surgical Nurse Manager.
ANESTHESIA PLAN (FOR GERBER HOSPITAL ONLY):
☐ General ☐ MAC
☐ Spinal
☐ Epidural
Regional block - Transfer of post-procedure pain management to the
Anesthesia Provider Surgeon concurs with the post-procedure Physician's Orders and
plan for intra-procedure anesthetic.

Telephone order/Verbal order documented and read-back completed. Practitioner's initials

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.



