

## Physician's Orders ORTHOPEDIC PRE-SCHEDULED SURGERY PRE-PROCEDURE

PRE-PROCEDURE	MRN Physician	
Page 1 of 2	rnysician	
Date of Surgery	FIN	
Surgeon/Physician		
Patient name		
REQUIRED: Prepare consent (Consent to read)		
REQUIRED: Weight kilograms (only) REQUIRED: Allerg	ies	
PENICILLIN ALLERGY? ☐ No ☐ Yes, reaction ☐ No anaphylaxis. May give Cephalosporin ☐ Anaphylaxis. No Cephalosporin		
REQUIRED (must choose one): ☐ A.M. Admit: Admit to Inpatient ☐ Outpatient ☐ Potential extended recovery (patient remains outpatient status, but may req ALL DESIRED ORDERS MUST BE CHECKED OR COMPLETED.	uire overnight stay. Final determination to be made post procedure)	
PHYSICIAN CONSULT:  Reason: Medical clearance. Name	LABORATORY:  All labs will be completed using Anesthesia Guidelines. Any orders checked below will be completed on Admission Day of Surgery, in addition to prior lab testing required by the Anesthesia Guidelines.   Laboratory studies/diagnostic tests Protocol - pre-procedure anesthesia orders: Pre-procedure - May initiate.  Basic metabolic panel	
Chlorhexidine cloth (CHG) skin cleansing pre-procedure Site  TEDs: ☐ Knee high ☐ Thigh high  Pneumatic compression device. Knee high bilaterally. Comments: For surgical cases lasting greater than 60 minutes and/or high risk patients as defined by Anesthesia Guidelines.  NOTE: If patient does not meet the Spectrum Health criteria above and you still want them placed, then SCD's must be ordered below. Pneumatic compression device: ☐ Knee high, bilaterally ☐ Knee high, right leg ☐ Knee high, left leg  Protocol - pre-procedure anesthesia orders: Pre-procedure - May initiate pre-operative anesthesia orders (include laboratory and diagnostic tests)	☐ TYPE AND SCREEN: PRBC's	

Patient Name

DOB

 ${\bf Telephone\ order/Verbal\ order\ documented\ and\ read-back\ completed.\ \ Practitioner's\ initials\ \_$ 

## **CONTINUED ON PAGE 2** →

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.

TRANSCRIBED:		VALIDATED:		ORDERED:		
TIME	DATE	TIME	DATE	TIME	DATE	Pager #
	Sign		R.N. Sign		Physician Print	Physician

CONFIDENTIAL NOTICE: The content of this fax is intended only for the named recipient(s) and may contain information that is protected under applicable law. If you are not the intended recipient(s) or if you receive this fax in error, notify the sender at the address or telephone number above. Also destroy any copies.

DO NOT MARK BELOW THIS LINE

BARCODE ZONE

DO NOT MARK BELOW THIS LINE





## **ORTHOPEDIC PRE-SCHEDULED SURGERY -**PRE-PROCEDURE (CONTINUED)

Page 2 of 2

ALL DESIRED	OPDERS MITST	RE CHECKED	OR COMPLETED.

IV SOLUTIONS:
1% lidocaine 0.25 to 2 mL ID for IV starts
Lactated ringers solution 1000 mL IV, 100 mL/hour
0.9% sodium chloride 1000 mL IV, 100 mL/hour
MEDICATIONS:
ANTIMICROBIALS (PROPHYLACTIC):  ☐ Cefazolin 2 grams IV, if patient is less than 120 kg administered per anesthesia ☐ Cefazolin 3 grams IV, if patient is greater than or equal to 120 kg administered per anesthesia
PENICILLIN (PCN) ALLERGY:
FOR IMMEDIATE TYPE PCN ALLERGY, CURRENT OR HISTORY OF MRSA COLONIZATION, MRSA RISK FACTORS:  [residence in long- term healthcare facility (not assisted living), HD patient, tracheostomy or gastrostomy tube, chronic decubitus ulcer or wound, long term vascular access]:
<ul> <li>Vancomycin (start administration within 120 minutes before skin incision):         <ul> <li>1 gram IV, if patient is less than 70 kg administered per anesthesia</li> <li>1.5 grams IV, if patient is 70-100 kg administered per anesthesia</li> <li>2 grams IV, if patient is greater than 100 kg administered per anesthesia</li> </ul> </li> </ul>
FOR IMMEDIATE TYPE PENICILLIN ALLERGY WHEN VANCOMYCIN IS NOT PREFERRED: ☐ Clindamycin 900 mg IV per anesthesia
OTHER MEDICATIONS:
<ul> <li>□ Tranexamic acid 1000 mg IV piggy back once routine, 10 minutes prior to incision, infuse over 10 minutes</li> <li>□ Tranexamic acid 1000 mg IV piggy back once routine, at tourniquet release, over 10 minutes</li> <li>□ Tranexamic acid 2000 mg IV piggy back once routine, single dose over 20 minutes prior to incision for cases without use of tourniquet</li> <li>□ Tranexamic acid 1950 mg PO once routine, 1-2 hours prior to incision</li> <li>□ Tranexamic acid 2000 mg/100 mL 0.9% sodium chloride irrigation topical, once routine. Wound soak for 5 minutes (for patients with risk or thrombosis with systemic tranexamic acid)</li> <li>□ Celecoxib 400 mg PO 1 hour pre-procedure</li> <li>□ Gabapentin 600 mg PO 1 hour pre-procedure</li> </ul>

Physician
MRN
DOB
Patient Name

	MRN	
	Physician	
	FIN	
MEDICATIO	ONS: (CONTINUED)	
RESPIRA	ATORY: ncentive spirometer	
ar th	OCKER: Continue current therapy with sips of vinesthesia for order if patient did not conerapy Medication	continue beta blocker
	Pose Route	
	For any additional orders: handv Must check the box for order to	be processed.
NOTE: •	If there is a frequent order that need contact Grand Rapids Spectrum He Planning Manager or Surgical Nurse	ealth Surgical Pre-procedure
☐ Gene ☐ Moni ☐ Spina ☐ Epidu ☐ Regic ☐ Anes ☐ Surge	itored anesthesia care (MAC) al	e pain management to the

Telephone order/Verbal order documented and read-back completed. Practitioner's initials \_

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.



CONFIDENTIAL NOTICE: The content of this fax is intended only for the named recipient(s) and may contain information that is protected under applicable law. If you are not the intended recipient(s) or if you receive this fax in error, notify the sender at the address or telephone number above. Also destroy any copies.