



Patient Name
DOB
MRN
Physician
CSN

Defaults for orders not otherwise specified below:

- Interval: Once
- Interval: PRN: Every _____ days
- Interval: Every _____ day

Duration:

- Until date: _____
- 1 year
- _____ # of Treatments

Anticipated Infusion Date _____ ICD 10 Code with Description _____

Height _____ (cm) Weight _____ (kg) Allergies _____

Site of Service

- SH Gerber
- SH Helen DeVos (GR)
- SH Lemmen Holton (GR)
- SH Ludington
- SH Pennock
- SH Reed City
- SH United Memorial
- SH Zeeland

Provider Specialty

- Allergy/Immunology
- Cardiology
- Gastroenterology
- Genetics
- Infectious Disease
- Internal Med/Family Practice
- Nephrology
- Neurology
- OB/GYN
- Other
- Otolaryngology
- Pulmonary
- Rheumatology
- Surgery
- Urology
- Wound Care

Appointment Requests

- For electrolyte replacement Appointments (Initial Order)
Status: Future, Expected: S, Expires: S+365, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after, Infusion
- For electrolyte replacement appointments (Subsequent Order) – FOR PRN ORDERS ONLY
Status: Future, Expected: S, Expires: S+365, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after, Infusion

Nursing Orders

- ONC NURSING COMMUNICATION 100
May Initiate IV Catheter Patency Adult Protocol

Electrolyte Replacement: Magnesium

- ONC SAFETY PARAMETERS AND SPECIAL INSTRUCTIONS 4
Magnesium serum level = 1.8 to 2.5 mg/dL. No intervention
- magnesium sulfate injection 2 g/50mL (Premix)
2 g, Intravenous, Administer over 60 Minutes (50 ml/hr), Once, Starting S, For 1 Doses
Magnesium serum level = 1.6 to 1.7 mg/dL. Total dose 2 Gm.
- magnesium sulfate injection 2 g/50mL (Premix)
2 g, Intravenous, Administer over 60 Minutes (50 ml/hr), Every 1 hour, Starting S, For 2 Doses
Magnesium serum level = 1.4 to 1.5 mg/dL. Total dose 4 Gm.
- magnesium sulfate injection 2 g/50mL (Premix)
2 g, Intravenous, Administer over 60 Minutes (50 ml/hr), Every 1 hour, Starting S, For 3 Doses
Magnesium serum level = 1.2 to 1.3 mg/dL. Total dose 6 Gm.
- magnesium sulfate injection 2 g/50mL (Premix)
2 g, Intravenous, Administer over 60 Minutes (50 ml/hr), Once, Starting S, For 1 Doses
Magnesium serum level LESS THAN 1.2 mg/dL: Initiate Magnesium Sulfate 2 Gm IVPB ONCE AND notify APP or attending for additional magnesium sulfate supplementation and/or laboratory orders

CONTINUED ON PAGE 2 →

NOTE: Epic Treatment/Therapy Plan Orders. To be scanned/attached to the appropriate Infusion Referral Order in Epic.

Confidentiality of this medical record shall be maintained except when use or disclosure is required or permitted by law, regulation, or written authorization by the patient.



Patient Name
DOB
MRN
Physician
CSN

Electrolyte Replacement: Potassium

- ONC SAFETY PARAMETERS AND SPECIAL INSTRUCTIONS 5**
Potassium serum level = 3.5 to 4.9 mmol/L: No intervention

- ONC SAFETY PARAMETERS AND SPECIAL INSTRUCTIONS 5**
Potassium GREATER THAN 5 mmol/L: Notify APP or attending AND release order for 12-lead electrocardiogram

- potassium chloride SA (K-DUR, KLOR-CON M) CR tablet 40 mEq**
40 mEq, Oral, Once, Starting S, For 1 Doses
Potassium serum level = 3.3 to 3.4 mmol/L. Give ORAL replacement if patient is able to tolerate PO.

- potassium chloride 20 mEq in 100mL bag (Premix)**
20 mEq, Intravenous, Administer 60 Minutes (100 ml/hr), Once, Starting S, For 1 Doses
Use IV route if patient is unable to tolerate oral replacement. Potassium serum level = 3.3 to 3.4 mmol/L. Total dose 20 mEq.

- potassium chloride 20 mEq in 100mL bag (Premix)**
20 mEq, Intravenous, Administer 60 Minutes (100 ml/hr), Every 1 hour, Starting S, For 2 Doses
Potassium serum level = 3.1 to 3.2 mmol/L. Total dose 40 mEq.

- potassium chloride 20 mEq in 100mL bag (Premix)**
20 mEq, Intravenous, Administer 60 Minutes (100 ml/hr), Every 1 hour, Starting S, For 3 Doses
Potassium serum level 2.8 to 3 mmol/L. Total dose 60 mEq. Check potassium level after completion of infusions.

- potassium chloride 20 mEq in 100mL bag (Premix)**
20 mEq, Intravenous, Administer 60 Minutes (100 ml/hr), Once, Starting S, For 1 Doses
For patients with serum potassium level LESS THAN 2.8 mmol/L initiate potassium chloride 20 mEq over 1 hour, notify provider and release order for 12-lead electrocardiogram; release order and collect sample for potassium blood level after completion of infusion.

- Electrocardiogram, Complete**
Routine, Hospital Performed, Status: Future, Expected: S, Expires: S+365

- Potassium, Blood Level**
Status: Future, Expires: S+365, URGENT, Clinic Collect, Blood, Blood, Venous

Confidentiality of this medical record shall be maintained except when use or disclosure is required or permitted by law, regulation, or written authorization by the patient.

Telephone order/Verbal order documented and read-back completed. Practitioner's initials _____

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.

TRANSCRIBED:		VALIDATED:		ORDERED:		Pager #
TIME	DATE	TIME	DATE	TIME	DATE	
	Sign		R.N. Sign		Physician Print	Physician