

## Corewell Health Surgical Services Hard Stops (HS)



Surgical Service Line	Surgeon	Procedure	HgbA1c Control	Tobacco/Nicotine Cessation	Decrease Opioid Use	Decrease BMI
Bariatric Surgery	Schram Foote Holman Smith Martin	All bariatric cases including revisions	HgbA1C >10%	Tobacco/Nicotine free (urine nicotine negative prior to surgery scheduled)	MME >60	No Upper Limit BMI, lower limit BMI ≥35
Colorectal Surgery	Figg Dujovny Hoedema Kim Luchtefeld Ogilvie Siripong Slay Gan	All elective cases (BMI specific to hernia and colostomy takedowns only)	HgbA1C >8.5%	Tobacco/Nicotine free (negative nicotine test 2-3 weeks from quit date)	MME >90	BMI ≥40 (ventral hernia and colostomy takedowns only)
ENT & ENT Oncology	Hart Postma Roche Wan Thompson	-Nasal cases -Elective mastoidectomies -Tympapanoplasties -Tonsillectomies	HgbA1C >8.5%	Tobacco/Nicotine free (negative nicotine test 2-3 weeks from quit date)	MME >90	NA
General Surgery	Banks-Venegoni Gawel Martin Minks Scheeres Silverman Smith Gallardo Conway Nava Durling	-Open complex ventral hernia repair with component separation -Laparoscopic/robotic complex ventral hernia repair with component separation -Open/Laparoscopic bowel resection	HgbA1C >7.5%	Tobacco/Nicotine free or counseling for elective cases (negative nicotine test 2-3 weeks from quit date)	MME >90	BMI ≥40 Laparoscopic hiatal hernia (type I) repair with fundoplication -Type II, III, IV paraesophageal hernias not applicable, Laparoscopic Nissen/Toupet/Dor fundoplication
Gynecology	Lalley Leary Platte Shah Friedman Sharma	-Hysterectomy -Myomectomy -Laparotomy -Laparoscopy -Prolapse surgery	HgbA1C >8.5%	Tobacco/Nicotine free Device and Mesh Procedures Only - Nicotine Free 8 weeks	MME >90	BMI ≥40
Gynecologic Oncology	Seamon Zakhour Hummel Gressel	Procedures as directed by the surgeon	HgbA1C >8.5%	Tobacco/Nicotine free (negative nicotine test 2-3 weeks from quit date)	MME >90	NA
Neurological Surgery	Boyce Patra Vitaz Mazaris Singer Ali Somji	Deformity procedures SI Joint fusions Elective lumbar fusions Elective cervical and lumbar fusions (at surgeon's discretion) <b>Spine exclusions:</b> microdiscs, laminectomies, spinal cord compression cases <b>Oncology exclusions:</b> neuro/oncology procedures	HgbA1C >8.0%	Tobacco/Nicotine free or counseling for elective cases, nicotine replacements are an option for spine cases. Urine nicotine testing is based on surgeon discretion	NA	NA
Orthopedic: Hand, Foot & Ankle	Buchanan Kelpin Hinkelman Chan Hess Duquette Buchanan Harris Stawiski Pangrazzi Erard	All elective hand, foot, and ankle procedures	HgbA1C >8.0%	Tobacco/Nicotine free (negative nicotine test 3-6 weeks from quit date)	MME >90	NA
Orthopedic - Spine	Googe Karek Squires Chu Brown Easton Kozlow Russo Stubbart Villerot	Deformity procedures SI Joint fusions Elective lumbar fusions Elective cervical and lumbar fusions (at surgeon's discretion) <b>Spine exclusions:</b> microdiscs, laminectomies, spinal cord compression cases	HgbA1C >8.0%	Tobacco free 2 weeks prior to surgery, nicotine replacements may be optional	NA	NA
Orthopedic Sports	Axtman Bouwkamp Hamilton Lebolt Menge Merillat	All elective Sports Medicine Procedures	HgbA1C >8.0%	Tobacco/Nicotine free Less than twice a week use (negative nicotine test 3-6 weeks from quit date)	MME >90	NA
Orthopedic - Total Joint	Sherry Perry Mahajan Ugolini Hood Grierson	All elective total joint procedures	HgbA1C >8.0%	Tobacco/Nicotine free (negative nicotine test 3-6 weeks from quit date)	MME >90	BMI ≥45
Ortho - Trauma	Conlan Moffitt George Maroto Starke	Procedures apply as directed by the surgeon	HgbA1C >8.0%	Tobacco/Nicotine Free No urine testing required <b>No NRTs should be prescribed</b>	MME >90	NA
Surgical Oncology	Assifi Chung Oostendorp Wright	Procedures apply as directed by the surgeon	HgbA1C >8.5%	Tobacco/Nicotine free or counseling for elective cases	MME >90	NA
Urology	Bolthouse Brede Hoekstra Humphrey Lane Pimentel Tobert Ghareeb Riedinger Southwell	Penile implants only	HgbA1C >8.5%	Tobacco free (negative nicotine test 2-3 weeks from quit date)	MME >90	BMI ≥40
Urologic Gynecology	Lalley Leary Platte Shah Friedman Sharma	All elective cases	HgbA1C >8.5%	Tobacco/Nicotine free Device and Mesh Procedures Only - Nicotine Free 8 weeks	MME >90	NA
Vascular	Bordoli Chambers Cuff Mansour Simmons Slaikau Kimball Kashyap Yassa	Claudicants for peripheral bypass surgery without wounds. 1st rib resection for neurogenic Thoracic Outlet Syndrome (TOS)	HgbA1C >8.5%	Tobacco free (negative nicotine test 2 weeks from quit date)	MME >90	NA

\*Optimize Malnutrition: Each patient is screened using MST Tool by SOC Nurse Navigator. The bariatric service screens its own patients for malnutrition.

\*Optimize Anemia: Each patient is screened for anemia (criteria Hgb &lt;13 for men and women)