

Corewell Health West Surgical Services Hard Stops (HS)



Surgical Service Line	Surgeon	Procedure	HgbA1c Control	Tobacco/Nicotine Cessation	Decrease Opioid Use	Decrease BMI
Bariatric Surgery	Schram - Foote - Holman - Smith - Martin	All bariatric cases including revisions	HgbA1C >10%	Tobacco/Nicotine free (urine nicotine negative prior to surgery scheduled)	MME >60	No Upper Limit BMI, lower limit BMI ≥35
Colorectal Surgery	Figg - Dujovny - Luchtefeld - Hoedema - Kim - Ogilvie - Siripong - Gan	All elective cases (BMI specific to hernia and colostomy takedowns only)	HgbA1C >8.5%	Tobacco/Nicotine free (negative nicotine test 2-3 weeks from quit date)	MME >90	BMI ≥40 (ventral hernia and colostomy takedowns only)
ENT & ENT Oncology	Hart - Postma - Roche - Wan - Thompson	-Nasal cases -Elective mastoidectomies -Tyumpanoplasties -Tonsillectomies	HgbA1C >8.5%	Tobacco/Nicotine free (negative nicotine test 2-3 weeks from quit date)	MME >90	NA
General Surgery	Autrey - Chapman - Dull - Durling - Fletter - Gawel - Haan - Kuiper - Kwong Hing - Martin - Price - Parsons - Pentiak - Rappaport - Schaefer - Smith - Spencer - VandenDool - Visser - Winkler - WRight - Yang - Zambito - Ziolkowski	-Open complex ventral hernia repair with component separation -Laparoscopic/robotic complex ventral hernia repair with component separation -Open/Laparoscopic bowel resection	HgbA1C >7.5%	Tobacco/Nicotine free or counseling for elective cases (negative nicotine test 2-3 weeks from quit date)	MME >90	BMI ≥40 Laparoscopic hiatal hernia (type I) repair with fundoplication -Type II, III, IV paraesophageal hernias not applicable, Laparoscopic Nissen/Toupet/Dor fundoplication
Gynecology	Lalley - Leary - Platte - Shah - Friedman-Sharma	-Hysterectomy -Myomectomy -Laparotomy -Laparoscopy -Prolapse surgery	HgbA1C >8.5%	Tobacco free Device and Mesh Procedures Only Prescription Nicotine Replacement Products may be used	MME >90	BMI ≥40
Gynecologic Oncology	Seamon - Zakhour - Hummel - Gressel	Procedures as directed by the surgeon	HgbA1C >8.5%	Tobacco/Nicotine free (negative nicotine test 2-3 weeks from quit date)	MME >90	NA
Neurological Surgery	Boyce - Clark - Figuero - Keller - Mazaris - Patra Singer - Somji - Vitaz - Vogel - Williams	Deformity procedures SI Joint fusions Elective lumbar fusions Elective cervical and lumbar fusions (at surgeon's discretion) <b>Spine exclusions:</b> microdiscs, laminectomies, spinal cord compression cases <b>Oncology exclusions:</b> neuro/oncology procedures	HgbA1C >8.0%	Tobacco/Nicotine free or counseling for elective cases, nicotine replacements are an option for spine cases. Urine nicotine testing is based on surgeon discretion	NA	NA
Orthopedic: Hand, Foot & Ankle	Buchanan - Chan - Duquette - Erard - Harris - Hess - Hinkelman - Kelpin - Pangrazzi - Stawiski	All elective hand, foot, and ankle procedures	HgbA1C >8.0%	Tobacco/Nicotine free (negative nicotine test 3-6 weeks from quit date)	MME >90	NA
Orthopedic - Spine	Chu - Brown - Easton - Googe - Kerek - Kozlow - Squires - Stubbart - Villerot	Deformity procedures SI Joint fusions Elective lumbar fusions Elective cervical and lumbar fusions (at surgeon's discretion) <b>Spine exclusions:</b> microdiscs, laminectomies, spinal cord compression cases	HgbA1C >8.0%	Tobacco free 2 weeks prior to surgery, nicotine replacements may be optional	NA	NA
Orthopedic Sports	Axtman - Bouwkamp - Hamilton - Lebolt - Menger - Merillat	All elective Sports Medicine Procedures	HgbA1C >8.0%	Tobacco/Nicotine free Less than twice a week use (negative nicotine test 3-6 weeks from quit date)	MME >90	NA
Orthopedic - Total Joint	Hood - Grierson - Hamilton - Mahajan - Perry - Sherry - Ugolini	All elective total joint procedures	HgbA1C >8.0%	Tobacco/Nicotine free (negative nicotine test 3-6 weeks from quit date)	MME >90	BMI ≥45
Ortho - Trauma	Conlan - George - Maroto - Moffitt - Starke	Procedures apply as directed by the surgeon	HgbA1C >8.0%	Tobacco/Nicotine Free urine testing required <b>No NRTs should be prescribed</b>	MME >90	NA
Surgical Oncology	Assifi - Chung - Oostendorp - Wright	Procedures apply as directed by the surgeon	HgbA1C >8.5%	Tobacco/Nicotine free or counseling for elective cases	MME >90	NA
Urology	Bolthouse - Brede - Curry - Drobish - Ghareeb - Hoekstra - Humphrey - Lane - Pimentel - Riedeinger - Rizer - Southwell - Tobert	Penile implants only	HgbA1C >8.5%	Tobacco free (negative nicotine test 2-3 weeks from quit date)	MME >90	BMI ≥40
Urologic Gynecology	Lalley Leary Platte Shah Friedman Sharma	All elective cases	HgbA1C >8.5%	Tobacco free Device and Mesh Procedures Only Prescription Nicotine Replacement Products may be used	MME >90	NA
Vascular	Bordoli Chambers Cuff Mansour Simmons Slaikeu Kimball Kashyap Yassa	Claudicants for peripheral bypass surgery without wounds. 1st rib resection for neurogenic Thoracic Outlet Syndrome (TOS)	HgbA1C >8.5%	Tobacco free (negative nicotine test 2 weeks from quit date)	MME >90	NA

\*Optimize Malnutrition: Each patient is screened using MST Tool by SOC Nurse Navigator. The bariatric service screens its own patients for malnutrition.

\*Optimize Anemia: Each patient is screened for anemia (criteria Hgb <13 for men and women)