

| <i>Short Name</i> | <i>Comments</i> | <i>Type</i> | <i>Investigator</i> | <i>Contact</i> |
|------------------------------|---|-------------|---------------------|--|
| Ketamine v. Lidocaine | A Prospective, Randomized, Single-Blinded Trial of Ketamine Versus Lidocaine Infusions for Multimodal Pain Management in Traumatic Rib Fracture Patients | Treatment | Brittany Hoyte | Breanna Barber breanna.barber@spectrumhealth.org 616.391.3329 |
| Sextant | This is a randomized clinical trial where participants who have suffered from open extremity fractures will either receive standard of care prophylactic antibiotic treatment (Vancomycin only) or study treatment (Vancomycin & Tobramycin) | Treatment | Dr. Moffitt | Breanna Barber breanna.barber@spectrumhealth.org 616.391.3329 |
| Tobra | This is a randomized clinical trial where participants who have suffered from tibula pilon or plateau will either receive standard of care prophylactic antibiotic treatment (Vancomycin only) or study treatment (Vancomycin & Tobramycin). | Treatment | Dr. Moffitt | Breanna Barber breanna.barber@spectrumhealth.org 616.391.3329 |
| Acute SCI Study | Phase 2, double-blinded, randomized, controlled clinical trial for acute spinal cord injury patients. MT-3921 (study drug) is a monoclonal antibody administered via IV infusion to help treat patients. | Treatment | Dr. Karek | Breanna Barber breanna.barber@spectrumhealth.org 616.391.3329 |