SPECTRUM HEALTH	Patient Name
Physician's Orders PERIPHERAL VASCULAR PRE-SCHEDULED SURGERY - PRE-PROCEDURE Page 1 of 2	DOB MRN Physician
Date of SurgerySurgeon/PhysicianPatient nameREQUIRED: Prepare consent (Consent to read)	Date of birth
REQUIRED: Weight kilograms (only)       REQUIRED: Aller         PENICILLIN ALLERGY?       No       Yes, reaction         No anaphylaxis. May give Cephalosporin       Anaphylaxis. No Cephalosporin         REQUIRED (must choose one):       A.M. Admit: Admit to Inpatient       Outpatient         Potential extended recovery (patient remains outpatient status, but may reconstruction of the construction	
<ul> <li>CONSULTS:</li> <li>PHYSICIAN CONSULT:</li> <li>Reason: Medical clearance. Name</li> <li>Reason: Consult surgical pain service for block</li> <li>PT CARE/ACTIVITY:</li> <li>Hair clipping pre-procedure in Surgical Center as indicated. Site</li> <li>Chlorhexidine cloth (CHG) skin cleansing pre-procedure Site</li> <li>Pneumatic compression device. Knee high bilaterally. Comments: For surgical cases lasting greater than 60 minutes and/or high risk patients as defined by Anesthesia Guidelines.</li> <li>NOTE: If your patient does not meet the Spectrum Health criteria above and you still want them placed, SCD's must be ordered below.</li> <li>Pneumatic Compression Device:</li> <li>Knee high, bilaterally</li> <li>Knee high, right leg</li> <li>Knee high, left leg</li> <li>Protocol - pre-procedure anesthesia orders: Pre-procedure - May initiate pre-operative anesthesia orders (include laboratory and diagnostic tests)</li> </ul>	LABORATORY:         All labs will be completed using Anesthesia Guidelines. Any orders checked below will be completed on Admission Day of Surgery, in addition to prior lab testing required by the Anesthesia Guidelines.         Basic metabolic panel       CMP         CBC with diff.       CBC without diff.         Protime (with INR)       PTT         Urinalysis (UA)       UA, culture if indicated         Lytes       Blood type, ABO/Rh typing         TYPE AND SCREEN: PRBC's number of units         Hemoglobin A1c level         POC pregnancy test urine (SH Grand Rapids)         Pregnancy qualitative urine (Other locations)         POC GLUCOSE TESTING:         For all patients with known diabetes         M Obtain Glucose POC once, then every 2 hours         NOTIFY:         Anesthesia, if blood glucose is greater than 180 or less than 70         DIAGNOSTICS:         Electrocardiogram (ECG)         IV SOLUTIONS:         M 1% lidocaine 0.25 to 2 mL ID for IV starts         Lactated ringers solution 1000 mL IV, 100 mL/hour         0.9% sodium chloride 1000 mL IV, 100 mL/hour

#### Telephone order/Verbal order documented and read-back completed. Practitioner's initials \_\_\_\_

# CONTINUED ON PAGE 2 →

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.

TRANSCRIBED:		VALIDATED:	VALIDATED:		ORDERED:		
TIME	DATE	TIME	DATE	TIME	DATE	Pager #	
	Sig	n	R.N. Sign		Physician Print		Physician

CONFIDENTIAL NOTICE: The content of this fax is intended only for the named recipient(s) and may contain information that is protected under applicable law. If you are not the intended recipient(s) or if you receive this fax in error, notify the sender at the address or telephone number above. Also destroy any copies.



+

 $^+$ 

 $^+$ 



# PERIPHERAL VASCULAR PRE-SCHEDULED SURGERY - PROCEDURE (CONTINUED)

Page 2 of 2

## ALL DESIRED ORDERS MUST BE CHECKED OR COMPLETED.

#### **MEDICATIONS:**

#### ANTIMICROBIALS (PROPHYLACTIC):

- Cefazolin 2 grams IV, if patient is less than 120 kg administered per anesthesia
- Cefazolin 3 grams IV, if patient is greater than or equal to 120 kg administered per anesthesia

## **PENICILLIN (PCN) ALLERGY:**

FOR IMMEDIATE TYPE PCN ALLERGY, CURRENT OR HISTORY OF MRSA COLONIZATION, MRSA RISK FACTORS:

[residence in long- term healthcare facility (not assisted living), HD patient, tracheostomy or gastrostomy tube, chronic decubitus ulcer or wound, long term vascular access]:

- Vancomycin (start administration within 120 minutes before skin incision):
  - 1 gram IV, if patient is less than 70 kg administered per anesthesia
  - □ 1.5 grams IV, if patient is 70-100 kg administered per anesthesia
  - $\Box$  2 grams IV, if patient is greater than 100 kg administered per anesthesia
- FOR IMMEDIATE TYPE PENICILLIN ALLERGY WHEN VANCOMYCIN IS NOT PREFERRED:
  - Clindamycin 900 mg IV per anesthesia

# **VTE PROPHYLAXIS (PHARMACOLOGIC:**

- Enoxaparin 40 mg subQ upon arrival **NOTE:** If spinal or epidural anesthesia, or peripheral nerve block are planned, DO NOT use enoxaparin. Use subQ heparin instead.
- Heparin 5000 unit(s) subQ upon arrival

Patient Name
DOB
MRN
Physician
FIN

#### MEDICATIONS: (CONTINUED)

## **RESPIRATORY:**

	Incentive	spirometer
--	-----------	------------

# BETA BLOCKER:

🗙 Continue current	therapy with sips of	water in AM. Contact
anesthesia for or	der if patient did not	continue beta blocker
therapy		
Medication		
Dose	Route	Frequency

Must check the box for order to be processed.

#### OTHER:

**NOTE:** • For any additional orders: handwrite clearly or type below.

NOTE: • If there is a frequent order that needs to be added to your form, contact Grand Rapids Spectrum Health Surgical Pre-procedure Planning Manager or Surgical Nurse Manager.

Telephone order/Verbal order documented and read-back completed. Practitioner's initials \_

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.

TRANSCRIBED:		VALIDATED:		ORDERED:			
TIME	DATE	TIME	DATE	TIME	DATE	Pager #	
	Sign		R.N. Sign		Physician Print	Physiciar	ı

CONFIDENTIAL NOTICE: The content of this fax is intended only for the named recipient(s) and may contain information that is protected under applicable law. If you are not the intended recipient(s) or if you receive this fax in error, notify the sender at the address or telephone number above. Also destroy any copies. X13425 (8/16) - Page 2 of 2

+