	Spectru Health
--	-------------------

Physician's Orders GOLIMUMAB (SIMPONI ARIA), IV PIGGYBACK -ADULT, OUTPATIENT, INFUSION CENTER Page 1 to 2

Patient Name	
DOB	
MRN	
Physician	
FIN	

Defaults for orders not otherwise specified below:

- □ Interval: INDUCTION Every 28 days x 2 treatments (maintenance treatment begins on day 84)
- □ Interval: MAINTENANCE Every 56 days

Duration:

- Until date: _____
- 1 year

of Treatments

Anticipated Infusion Da	te ICD 10	Code with Des		
Height	(cm) Weight	(kg) Allergies_		
Site of Service				
SH Gerber	□ SH Lemmen Holton (G	R)	SH Pennock	SH United Memorial
□ SH Helen DeVos (GR)	SH Ludington		SH Reed City	SH Zeeland
Provider Specialty				
□ Allergy/Immunology	Infectious Disease		□ OB/GYN	Rheumatology
Cardiology	□ Internal Med/Family Pra	actice	□ Other	□ Surgery
Gastroenterology	Nephrology		Otolaryngology	Urology
Genetics	Neurology		Pulmonary	□ Wound Care

Appointment Requests

Infusion Appointment Request Status: Future, Expected: S, Expires: S+365, Sched. Tolerance: Schedule appointment at most 3 days before or at most 3 days after, Infusion and possible labs

Safety Parameters and Special Instructions

~	ONC SAFETY PARAMETERS AND SPECIAL INSTRUCTIONS 4
	GOLIMUMAB (SIMPONI ARIA)

An FDA-approved patient medication guide, which is available with the product information and as follows, should be dispensed with this medication

Https://www.accessdata.fda.gov/drugsatfda_docs/label/2017/125433s019lbl.pdf

Treatment with SIMPONI ARIA should not be initiated in patients with an active infection, including clinically important localized infections.

Tuberculosis surveillance and maintenance: Screen and treat latent infection prior to starting therapy. Hepatitis B surveillance and maintenance: Screen prior to initiating therapy. Refer to specialist as warranted by serology.

TB skin test, hepatitis B surface antigen (HBsAg) test, liver function test (LFT), complete blood count (CBC), up-to-date vaccinations, risk assessment for cancer, and pregnancy testing. Monitor for signs of tuberculosis throughout therapy. Do not initiate therapy if active infection is present. Monitor closely for signs and symptoms of infection. Monitor for signs/symptoms of malignancy (eg, splenomegaly, hepatomegaly, abdominal pain, persistent fever, night sweats, weight loss). Identify history of latex or polysorbate 80 allergy; some dosage containers may contain these agents. Monitor LFTs, CBC at regular intervals. Assess results of laboratory tests (PDD) at regular intervals during treatment.

	ONC SAFETY PARAMETERS AND SPECIAL INSTRUCTIONS 6
	Verify all INDUCTION/LOADING DOSES given prior to start of MAINTENANCE DOSES
	ONC SAFETY PARAMETERS AND SPECIAL INSTRUCTIONS 4
_	HEPATITIS B VIRUS SURVEILLANCE AND MAINTENANCE RECOMMENDATIONS: Screen prior to treatment. Refer to specialist as warranted by serology.

ONC SAFETY PARAMETERS AND SPECIAL INSTRUCTIONS 5 TUBERCULOSIS SURVEILLANCE AND MANAGEMENT RECOMMENDATIONS: Screen prior to treatment. Treat latent infection prior to starting therapy.

CONTINUED ON PAGE 2 →

NOTE: Epic Treatment/Therapy Plan Orders. To be scanned/attached to the appropriate Infusion Referral Order in Epic.

+

+



Spectrum GOLIMUMAB Health (SIMPONI ARIA), IV PIGGYBACK -ADULT, OUTPATIENT, INFUSION CENTER (CONTINUED) Page 2 to 2

Patient Name	
DOB	
MRN	
Physician	

FIN

Labs Interval Duration Complete Blood Count w/Differential ✓ Status: Future, Expected: S, Expires: S+184, URGENT, Clinic Collect, Blood, Blood, Venous Hepatic Function Panel (Liver Panel) \checkmark Status: Future, Expected: S, Expires: S+365, URGENT, Clinic Collect, Blood, Blood, Venous Hepatitis B Surface Antigen Level Once 1 treatment Status: Future, Expected: S, Expires: S+365, URGENT, Clinic Collect, Blood, Blood, Venous Hepatitis B Core Total Antibody Level Once 1 treatment Status: Future, Expected: S, Expires: S+365, URGENT, Clinic Collect, Blood, Blood, Venous Arrange For Patient To Have Id Tb Skin Test Administered And Read Or Serum Tb Screening Lab Prior To Therapy Or Annually **ONC PROVIDER REMINDER 28** Once 1 treatment Arrange for patient to have intradermal TB skin test (tuberculin PPD) screening performed and read prior to initiating therapy and annually. TB Screen (Quantiferon Gold) Once 1 treatment Status: Future, Expected: S, Expires: S+365, URGENT, Clinic Collect, Blood, Blood, Venous Additional Lab Orders Labs: Every days Until date: Once 1 year # of Treatments **Nursing Orders** ONC NURSING COMMUNICATION 100 May Initiate IV Catheter Patency Adult Protocol **Treatment Parameters** ~ **ONC MONITORING AND HOLD PARAMETERS 3** May proceed with treatment if hepatitis B core antibody and surface antigen labs have been resulted prior to the first dose, and the results are negative. **ONC MONITORING AND HOLD PARAMETERS 4** \checkmark May proceed with treatment if tuberculosis screening test with either TB Screen blood test (QuantiFERON® Gold Plus) or TB skin test have been resulted prior to first dose and the results are negative. Medications golimumab (SIMPONI ARIA) 2 mg/kg in sodium chloride 0.9 % 100 mL IVPB **|** 2 mg/kg, Intravenous, Administer over 30 Minutes, Once, Starting S, For 1 Dose Infuse over 30 minutes. Do not infuse in the same line with other medications.

Telephone order/Verbal order documented and read-back completed. Practitioner's initials _____

NOTE: Unless Order is written DAW (dispense as written), medication may be supplied which is a generic equivalent by nonproprietary name.

ſ	TRANSCRIBED:		VALIDATED:		ORDERED:		
	TIME	DATE	TIME	DATE	TIME	DATE	Pager #
		Sign		R.N. Sign		Physician Print	Physician

EPIC VERSION DATE: 09/12/20

X25215 (7/21) - Page 2 of 2 © Spectrum Health

+