


Decision-making and organisation of reperfusion strategies within first 12 hours of medical contact



Note: Adapted from Chew et al.

Abbreviations: EMS, emergency medical service; FMC, first medical contact; PCI, percutaneous coronary intervention; STEMI, ST-segment elevation myocardial infarction.

Reference

Chew DP, Scott IA, Cullen L, et al. National Heart Foundation of Australia and Cardiac Society of Australia and New Zealand: Australian clinical guidelines for the management of acute coronary syndromes 2016. Medical Journal of Australia. 2016;205:128-33.

Brieger D, Cullen L, Briffa T et al. National Heart Foundation of Australia & Cardiac Society of Australia and New Zealand: Comprehensive Australian clinical guideline for diagnosing and managing acute coronary syndromes 2025. 2025. <https://www.heartfoundation.org.au/for-professionals/acs-guideline>.

This article was adapted from Heart, Lung and Circulation, 34(4), Brieger D, Cullen L, Briffa T et al., National Heart Foundation of Australia & Cardiac Society of Australia and New Zealand: Australian clinical guideline for diagnosing and managing acute coronary syndromes 2025, 309-397, Copyright Elsevier (2025). <https://doi.org/10.1016/j.hlc.2025.02.102>.

HH-PRISK-096.1.0425

View the **Acute coronary syndromes (ACS)** guideline

