

Made Possible.

Johnson Johnson vision

TECNIS Synergy™IOL



Powered by InteliLight™

About TECNIS Synergy<sup>™</sup> and TECNIS Synergy<sup>™</sup> Toric IOL Physicians considering use of the TECNIS Synergy<sup>™</sup> and TECNIS Synergy<sup>™</sup> Toric IOLs should refer to the Directions for Use labeling for a complete list of indications and safety information. See reverse for continued Indications and Important Safety Information.

#### What is a cataract?

Cataracts form as we age and are caused by a buildup of the eye's natural protein. The increase in protein causes a cloudiness in the lens of the eye, which is why people with cataracts may experience blurry or dull vision.<sup>10</sup>

### Do you have presbyopia?

Presbyopia typically occurs around age 40 and is an age-related loss of the eyes' ability to see objects up close. Cataract treatment is an opportunity to correct presbyopia the time of surgery, with a lens like **TECNIS Synergy™ IOL**, which offers a continuous range of vision from up close to far away, day and night. 347,911

### Do you have astigmatism?

Astigmatism is when the shape of your eye is more oblong than circular, which can create distorted vision.
With TECNIS Synergy™ IOL TORIC II astigmatism can be corrected during cataract surgery resulting in less need for glasses.™ Talk to your doctor about your astigmatism correction options.

### TECNIS Synergy<sup>™</sup> IOL A Presbyopia-Correcting Lens





Images are an illustrative simulation and do not represent actual data derived from studies.

## Do more of what you love, day and night

Cataract surgery is your opportunity to improve your vision from cloudy or blurry to crisp and clear.

At Johnson & Johnson Vision, we believe that clear vision shouldn't be limited to daylight and we strive to improve quality of life after cataract surgery, both day and night.

Think about walking downstairs in dim light, or seeing objects in the road when driving at night. These activities can be troublesome, which is why we developed InteliLight."





# Bring on the Night with InteliLight™

Contrast sensitivity is your ability to recognize an object against its background. It is important, especially in low light, because it allows you to see things like furniture in the dark, objects while driving at night, or even a curb at the end of the sidewalk when walking in the evening.

InteliLight<sup>™</sup> lens technology is the solution and offers the best contrast sensitivity and low light vision<sup>0</sup>, to help you see clearly day and night, even in dim light.<sup>2,3,5,6</sup>


## The widest range of continuous vision of

**TECNIS Synergy™ IOL** is a premium intraocular lens (IOL) used to treat cataracts.

It gives you the best near vision,\* while also helping you to see clearly across the widest range of vision — near, intermediate, and far. This lens is powered by our InteliLight™ technology, so that you can see better both day and night.<sup>3-7</sup>

Learn more at:

www.tecnisvisionsimulator.com



### Confidence in any light



Fewer visual disturbances when driving at night.<sup>5</sup>



Freedom to do more after sunset. 49



Reduction in the risk of falls. 2,3,5,6,12,13

### TECNIS Synergy\*IOL



Toric II

INDICATIONS AND IMPORTANT SAFETY INFORMATION FOR TECNIS SYNERGY"
AND TECNIS SYNERGY" TORIC II IOL WITH TECNIS SIMPLICITY DELIVERY SYSTEM,
MODELS DEROOV AND DEW

CAUTION: Federal law restricts this device to sale, distribution and use by or on the order of a physician.

INDICATIONS FOR USE: The TECNIS Simplicity\* Delivery System is used to fold and assist in inserting the TECNIS Synergy\* IOL, which is indicated for primary implantation for the visual correction of aphakia in adult patients, with less than 1 diopter of pre-existing corneal astigmatism, in whom a cataractous lens has been removed. The TECNIS Simplicity\* Delivery System is used to fold and assist in inserting the TECNIS Synergy\* Toric II IOLs, which are indicated for primary implantation for the visual correction of aphakia and for reduction of refractive astigmatism in adult patients, with greater than or equal to 1 diopter of preoperative corneal astigmatism, in whom a cataractous lens has been removed. Compared to an aspheric monofocal lens, the TECNIS Synergy\* IOLs mitigate the effects of presbyopia by providing improved visual acuity at intermediate and near distances to reduce eyeglass wear, while maintaining comparable distance visual acuity. The lens is intended for capsular bag placement only.

#### CONTRAINDICATIONS: None.

RISKS: Routine cataract surgery risk, irrespective of lens selection, could be minor, temporary, or affect patients' vision permanently. Rare complications are worsening of vision, bleeding, or infection. Risks related to use of this lens include halo or glare in nighttime or poor visibility conditions. These may be perceived as an annoyance or hindrance, which, on rare occasions, may be significant enough for the patient to request removal of the IOL. Patients should discuss all risks and benefits with their eye doctor before surgery.

PRECAUTIONS: Recent contact lens usage may affect the patient's refraction; therefore, in contact lens wearers, surgeons should establish corneal stability without contact lenses prior to determining IOL power. Prior to surgery, the surgeon must inform prospective patients of the possible risks and benefits associated with the use of this device and provide a copy of the patient information brochuse to the patient. Patients with a predicted postoperative astigmatism greater than 1.0 D may not be suitable candidates for implantation with the TECNIS Synergy" and TECNIS Synergy" Toric IOLs, as they may not obtain the benefits of reduced spectacle wear or improved intermediate and near vision seen in patients with lower postoperative astigmatism. The safety and effectiveness of the TECNIS Synergy" IOL have not been substantiated in patients under the age of 22 or those with preexisting ocular conditions and intraoperative complications.

WARNINGS: Physicians considering lens implantation should weigh the potential risk/benefit ratio. Well-informed patients with well-defined visual needs and preferences should be selected for lens implantation. Patients should be informed of the possibility of visual effects (such as halo or glare around lights), which may be expected in nighttime or poor visibility conditions.

- \* vs. AcrySof\* IQ PanOptix\*, TECNIS Symfony™, TECNIS\* Multifocal. Based on comparison of DFU defocus curves and head to head clinical study vs. PanOptix\*.
- \*\*\* Individual results will vary. Some TECNIS Synergy™ patients may require spectacles post-surgery.
- † Continuous 20/32 or better.
- ‡ Based on interim data collected at 6 months post operative, 92% of subjects did not wear glasses.
- ◊ vs. PanOptix® IOL.

#### REFERENCES:

- DOF2020CT4011: DOF\_Effect of blocking violet light on light scatter in Tecnis Symfony™ IOLs (v1.0).
- 2. Data on file. DOF2020OTH4010 Johnson & Johnson Vision, Inc. Santa Ana, CA.
- Data on file. Johnson & Johnson Surgical Vision, Inc. DOF2019OTH4002 Weeber H. MTF of the TECNIS Synergy™ OptiBlue¹ IOL, and other lens models. 27 Mar 2019.
- TECNIS Symfony™ Extended Range of Vision IOL DFU US -Doc. #z311215. Rev. 01, 12/2017 REF2020MLT4051.
- 5. Data on file. DOF2020OTH4011 Johnson & Johnson Vision, Inc. Santa Ana, Calif.
- 6. Data on file. DOF2015CT0020 Johnson & Johnson Vision, Inc. Santa Ana, Calif.
- TECNIS<sup>®</sup> Synergy<sup>™</sup> IOL with TECNIS Simplicity<sup>®</sup> Delivery System DFU, Z311421E.
- Presbyopia. American Optometric Association Website: https://www.aoa.org/patients-and-public eye-and-vision-problems/glossary-of-eye-and-visionconditions/presbyopia.
- DOF2020CT4015 Forte 1: A Comparative Clinical Evaluation of a New TECNIS' Presbyopia Correcting Intraocular Lens Against a PanOptix' Intraocular Lens- SPECTACLE WEAR AND SATISFACTION RESULTS.
- National Eye Institute. Cataracts. https://www.nei.nih.gov/learn-about-eye-health/eye-conditionsand-diseas... Accessed May 24, 2022.
- TECNIS' Multifocal 1-Piece IOL ZKB00 and ZLB00 DFU US Doc. #Z311328. Rev. A, 04/2018. REF2019CT4049.
- Rosenthal B, Fischer M. (2014) Functional vision changes in the normal and aging eye. In TL Kauffman (Ed.) Geriatric Rehabilitation Manual (2nd Ed, Chpt 51, pg 381-391). China: Elsevier Ltd.
- 13. Marks R. Falls among the elderly and vision: A Narrative Review. Open Med J 2014;1:54-65.

