


Drug Safety Overview: Pregnancy



Providers: Review the following lists of medications that are compatible with pregnancy, and those that are NOT compatible with pregnancy.



Pregnancy compatible

- Azathioprine
- Chloroquine
- Colchicine
- Cyclosporine
- Hydroxychloroquine
- NSAIDs (discontinue at 20 weeks)
- Prednisone (<20 mg/day)
- Sulfasalazine
- Tacrolimus



NOT compatible with pregnancy

MEDICATIONS

NOTES

Belimumab	Discuss discontinuation at conception
Cyclophosphamide	Consider if life/organ threatening disease in 2nd or 3rd trimester
Leflunomide	Use cholestyramine washout until level is undetectable
Methotrexate	Stop 1-3 months before trying to conceive
Mycophenolate mofetil; Mycophenolic acid	Discontinue 6 weeks before trying to conceive
Rituximab	Can be continued until conception; Consider if life/organ threatening disease in 2nd or 3rd trimester
Thalidomide; Lenalidomide	Stop 1 month before trying to conceive

These educational materials were supported by a medical education grant from GlaxoSmithKline to address Systemic Lupus Erythematosus (SLE) and Reproductive Health education to support patients, educators, and clinical teams. Its contents are solely the responsibility of the authors and do not necessarily reflect the official views of GlaxoSmithKline.

© 2021 American College of Rheumatology. All rights reserved.