
PHARMACOLOGICAL THERAPY



88 studies included:

NSAIDs -

PICO 1: 1 RCT; PICO 33: 1 observational study; PICO 2: 5 RCTs;

PICO 34: no studies; PICO 3: 20 RCTs; PICO 35:no studies

Glucocorticoids -

PICO 4: 1 RCT, 3 observational studies; PICO 36: no studies;

PICO 13: 2 RCTs, 2 observational studies; PICO 45: no studies;

PICO 14: no studies; PICO 46: no studies; PICO 15: no studies:

PICO 47: no studies

Tumor Necrosis Factor Inhibitors (TNFi) and non-TNFi Biologics -

PICO 5: 4 RCTs, 1 observational studies; PICO 37: no studies;

PICO 6: 13 RCTs; PICO 38: 5 RCTs, 2 observational studies;

PICO 8: 1 RCT, 5 observational studies; PICO 40: no studies;

PICO 9: 3 observational studies; PICO 41: no studies;

PICO 10: no studies; PICO 42: no studies

Slow-Acting Anti-Rheumatic Drugs -


PICO 7: 15 RCTs; PICO 39: 1 RCT, 1 observational studies

Treatment of Stable Disease -


PICO 11: no studies; PICO 43: no studies;


PICO 12: 1 observational studies; PICO 44: no studies

REHABILITATION/PHYSICAL THERAPY




SURGICAL TREATMENT






INFLAMMATORY BOWEL DISEASE



PREVENTIVE CARE



DISEASE ACTIVITY MEASURES

